APPROACHES FOR ASSESSING HOMELESS CONTRIBUTIONS TO BACTERIAL CONTAMINATION

Presentation to the Water Quality Coordinating Committee

Stephen B. Weisberg

Southern California Coastal Water Research Project Authority

October 23, 2018

BACKGROUND

- Bacterial water quality impairment from homeless fecal deposits is a potential problem
 - Orange County Public Works removed 5,279 lbs of human fecal matter when they conducted their Santa Ana River cleanup effort
 - Human fecal material contains more pathogens of concern than other sources
- There is a long way from "potential" to "problem"
 - Don't know how much homeless contributions end up in the river
 - There have been no definitive studies quantifying bacterial load from homeless
- The scientific tools to help quantify the problem are evolving

THE CHANGING MANAGEMENT QUESTION


- Initial question was: "Is there a bacterial problem"?
- Next question: "How much of our bacterial problem is attributable to human fecal sources?"
- Question now: "How much of the human fecal contribution can be attributed to different sources?"
 - Private sewer laterals and septics
 - Public sewerage conveyance systems
 - Homeless
- Scientists are working on new techniques that will help answer the last question

IS THERE A BACTERIAL PROBLEM?

- Enterococcus and fecal coliforms are your core indicators
 - Not pathogenic themselves, but occur abundantly in the human gut
 - Epidemiological studies have shown their presence correlates with human illness
 - These are incorporated in your recently adopted bacterial provisions
- In wet weather, these indicator bacteria widely exceed objectives, regardless of location
- However, these indicators are of limited value for assessing homeless contributions
 - They don't differentiate human from animal fecal material
 - Could potentially do upstream-downstream studies, but they haven't been done
 - Could potential do a mass loading comparison, but many assumptions

HOW MUCH IS ATTRIBUTABLE TO HUMAN SOURCES?

- Next step Scientists identified a series of genetic markers that differentiate human from non-human sources (HF183)
 - Also identified markers for dog, gull, cow and horse feces
- Enhances opportunity for identifying potential fecal sources
 - Tremendous boon to TMDL planning
 - Also useful for prioritizing which locations to clean
- However, we frequently find human fecal markers in stormwater systems
 - HF183 is of no help in differentiating among human sources



California Microbial Source Identification Manual

COMMUNITY PROFILING

- New approach is microbial community profiling
 - Don't look for a specific genetic marker
 - Instead look for a community signature

Based on the concept that a unique microbial community grows

on biofilms inside sewer pipes

 If sewer microbial community found in runoff, then sewer contributions likely


 If not found in runoff, then another human source is responsible

STUDIES ARE UNDERWAY

- Sample many locations to evaluate spatial consistency
 - Sewerage system with different source inputs (e.g. residential vs. industrial)
 - Age of pipe
 - Pipe construction materials
- Sample over time to evaluate temporal consistency
 - Seasonality
 - Biofilm regrowth
- Initial results are extremely promising!




Preliminary Results: Microbial Community Analysis



MOVING FORWARD

- Studies this winter to further confirm that microbial communities differ between sanitary sewer and storm drains
 - We will be quantifying this at about 60 sites over the next year
- Will also be studying persistence and sensitivity
 - Not much value if the signal degrades quickly
 - Also not of much value if we can't measure it after dilution.
- San Diego as a case study for how to use the outcome
 - Study is a collaboration among the Regional Board, City and County
 - Not yet focused on developing similar signatures for homeless or septic
 - However, will be a big advance if we can confirm or eliminate sewer conveyance system leakage as a dominant source

Initial Results: Non-targeted Chemistry



DATA NECESSARY FOR ESTIMATING HOMELESS CONTRIBUTIONS

- Number of homeless
 - Locations and times

- Quantify the sanitary habits
 - Directly into the river
 - Indirectly into the river (upland or on riverbanks)
 - Offsite services provided by the cities
- Washoff experiments for upland or riverbank deposits